

# Immunoglobulin order form

Infusion Pharmacy Phone: 1-877-342-9352 Fax: 1-888-594-4844

Page 1 of 2

**IG specialist:** First Name: \_\_\_\_\_ Middle: \_\_\_\_\_ Last: \_\_\_\_\_ Phone: \_\_\_\_\_

**Patient information**  see attached  PEDIATRIC (younger than 13 years or less than 45kg in weight).

Patient First Name: \_\_\_\_\_ Middle: \_\_\_\_\_ Last: \_\_\_\_\_ Gender:  M  F DOB: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ ZIP: \_\_\_\_\_

Phone: \_\_\_\_\_  Home  Work  Cell Phone: \_\_\_\_\_  Home  Work  Cell

Emergency contact: \_\_\_\_\_ Phone: \_\_\_\_\_ Relationship: \_\_\_\_\_

**Insurance:**  Front and back of insurance cards attached.

Primary Insurance: \_\_\_\_\_ Phone: \_\_\_\_\_ Policy #: \_\_\_\_\_ Group: \_\_\_\_\_

Secondary Insurance: \_\_\_\_\_ Phone: \_\_\_\_\_ Policy #: \_\_\_\_\_ Group: \_\_\_\_\_

## Medical assessment

**Primary diagnosis ICD-10 code (required):** \_\_\_\_\_

Height in inches: \_\_\_\_\_ Weight **in kg only:** \_\_\_\_\_ Date weight (in kg) obtained: \_\_\_\_\_

Current medications?  Yes  No If yes, list here or attach a list: \_\_\_\_\_

Allergies: \_\_\_\_\_

Patient requires a first lifetime dose and is to receive the first dose in the home or Ambulatory Infusion Suite.

## Prescription and orders Medication to be infused per drug prescribing information recommended rate via a rate controlled device.

**Immune Globulin:**  No preference  Preferred product: \_\_\_\_\_ Dose will be rounded to the nearest vial or prefilled syringe size available.

**Directions:**  Infuse IV  Infuse SC  Titrate per manufacturer guidelines or as written: \_\_\_\_\_

**Initial loading:** \_\_\_\_\_ gm/kg divided over \_\_\_\_\_ days every \_\_\_\_\_ weeks; OR \_\_\_\_\_ gm/day x \_\_\_\_\_ days every \_\_\_\_\_ weeks.

**Maintenance:** \_\_\_\_\_ gm/kg divided over \_\_\_\_\_ days every \_\_\_\_\_ weeks; OR \_\_\_\_\_ gm/day x \_\_\_\_\_ days every \_\_\_\_\_ weeks.

**Other:** \_\_\_\_\_

**Quantity/Refills:** 1-month supply; refill x 1 year unless otherwise noted. Other: \_\_\_\_\_

Pharmacy to dispense flushes, needles, syringes and HME/DME in quantity sufficient to complete therapy as prescribed.

**Premedication:** Dispense PRN x 1 year (select below):

	Drug	Patient Type	Dose	Dispense detail	Directions
<input type="checkbox"/>	DiphenhydrAMINE	Adult & Pediatric > 30 kg	50 mg (two 25 mg capsules or tablets)	Dispense 25 mg capsules or tablets #100	Administer orally 30 minutes prior to Ig therapy. May repeat once if symptoms occur.
		Pediatric 15 - 30 kg	25 mg (10 mL)	Dispense 2.5 mg / mL oral solution 120 mL	
		Pediatric < 15 kg	12.5 mg (5 mL)	Dispense 2.5 mg / mL oral solution 120 mL	
<input type="checkbox"/>	Acetaminophen	Adult & Pediatric > 30 kg	325 mg	Dispense 325 mg tablets or 325 mg (10.15 mL) unit dose oral solution #100	Administer orally 30 minutes prior to Ig therapy. May repeat once if symptoms occur.
		Pediatric 15 - 30 kg	160 mg (5 mL)	Dispense 160 mg tablets #100 or 32 mg / mL oral solution 120 mL	
		Pediatric < 15 kg	80mg (2.5mL)	Dispense 32 mg / mL oral solution 120 mL	
<input type="checkbox"/>	Hydration - Sodium Chloride 0.9% <b>specify volume and rate</b>	Adult & Pediatric	Volume _____ mL	Dispense bag(s) for infusion #QS	Infuse IV prior to IG, at a rate of: <input type="checkbox"/> up to 250 mL / hr <input type="checkbox"/> up to 500 mL / hr <input type="checkbox"/> up to 900 mL / hr
<input type="checkbox"/>	Lidocaine-Prilocaine Cream 2.5%	SCIG & Pediatric	n/a	Dispense 30 Gm	Prior to administration of IG apply pea size amount topically to needle site(s).
<input type="checkbox"/>	Other, specify _____	_____	_____	_____	_____

**Lab Draw Orders x1 year (specify):** CMP  monthly  other \_\_\_\_\_ Serum creatinine/BUN  monthly  other \_\_\_\_\_

**Other lab (specify):** \_\_\_\_\_ Frequency  once  monthly  other \_\_\_\_\_

Lab work to be obtained via IV access using aseptic technique. If RN is not able to draw labs from a central catheter, the labs may be drawn peripherally. RN to flush IV access after each blood draw with 20 mL of 0.9% Sodium Chloride. As final lock for patency, RN to use 5 mL of heparin 10 units / mL. If therapy is being administered through an implanted port, use 5 mL of heparin 100 units / mL.

Please fax both pages of this completed form with a copy of any medical history and labs relevant to the prescribed therapy.

**This form is not a valid prescription in New York.**

# Immunoglobulin order form

Infusion Pharmacy Phone: 1-877-342-9352 Fax: 1-888-594-4844

Page 2 of 2

Patient First Name: \_\_\_\_\_ Middle: \_\_\_\_\_ Last: \_\_\_\_\_ DOB: \_\_\_\_\_

## Nursing orders

RN to complete assessment and administer IVIG via ambulatory pump or teach SCIG self-administration via appropriate pump (e.g., syringe, ambulatory), in the home or Ambulatory Infusion Suite. RN to insert / maintain / remove peripheral IV (PIVC) or access central venous catheter as needed using aseptic technique. RN to flush catheter with 5 mL of 0.9% Sodium Chloride pre infusion and post infusion. RN to rotate PIVC as needed for signs of infiltration or irritation.

If port, RN to access with non-coring port needle using sterile technique. De-access after infusion and apply pressure with sterile gauze. Apply transparent dressing to site. RN to use 10 mL sterile field 0.9% Sodium Chloride with needle change. Flush port with 10 mL of 0.9% Sodium Chloride pre infusion and post infusion. To maintain line patency following the post infusion flush, use 5 mL of heparin 100 units / mL. Discontinue port maintenance upon discontinuation of pharmacy services.

**Anaphylaxis/infusion reaction management orders:** Dispense PRN x 1 year

Therapy Type	Drug	Patient Type	Dose	Dispense detail	Directions*
IVIG	DiphenhydrAMINE (for mild to severe symptoms)	Adult & Pediatric > 30 kg	50 mg (two 25 mg capsules or tablets)	Dispense 25 mg capsules or tablets #4	For mild* symptoms, RN to slow infusion rate by 50% until symptoms resolve. Administer diphenhydrAMINE orally once. May repeat once if symptoms persist.
			50 mg (1 mL)	Dispense 50 mg / mL, 1 mL vial for injection #1	
		Pediatric 15 - 30 kg	25 mg (10 mL)	Dispense 2.5 mg / mL oral solution 120 mL (300 mg)	
			25 mg (0.5 mL)	Dispense 50 mg / mL, 1 mL vial for injection #1	
		Pediatric < 15 kg	12.5 mg (5 mL)	Dispense 2.5 mg / mL oral solution 120 mL (300 mg)	
			12.5 mg (0.25 mL)	Dispense 50 mg / mL, 1 mL vial for injection #1	
IVIG	EPINEPHrine (for severe symptoms)	Adult & Pediatric > 30 kg	0.3 mg (0.3 mL)	Dispense 1 mg vial for injection #2	For severe* symptoms (anaphylaxis), stop infusion. Disconnect tubing from access device to prevent further administration.
		Pediatric 15 - 30 kg	0.15 mg (0.15 mL)	Dispense 1 mg vial for injection #2	
		Pediatric 7.5 kg - 15 kg	0.1 mg (0.1 mL)	Dispense Autoinjector Pen 0.1 mg (PED) #2	Activate 911. Administer EPINEPHrine as an IM injection into the lateral thigh. Repeat EPINEPHrine in 5 to 15 minutes if symptoms persist. Initiate 0.9% Sodium Chloride IV. Administer CPR if needed until EMS arrives. Contact prescriber to communicate patient status.
SCIG	EPINEPHrine (for severe symptoms)	Adult & Pediatric > 30 kg	0.3 mg (0.3 mL)	Dispense Autoinjector Pen 0.3 mg #2	
		Pediatric 15 - 30 kg	0.15 mg (0.15 mL)	Dispense Autoinjector Pen JR 0.15 mg #2	
		Pediatric 7.5 - 15 kg	0.1 mg (0.1 mL)	Dispense Autoinjector Pen 0.1 mg (PED) #2	
IVIG	0.9% Sodium chloride (for severe symptoms)	Adult & Pediatric	500 mL	Dispense 500 mL bag #1	For severe symptoms administer as IV gravity bolus (1000 mL / hour).
IVIG	Other, specify _____	_____	_____	_____	_____

\*Mild symptoms include flushing, dizziness, headache, apprehension, sweating, palpitations, nausea, pruritus, and / or throat itching.

Moderate symptoms include chest tightness, shortness of breath, >20 mmHg change in systolic blood pressure from baseline, and / or increase in temperature (>2°F).

Severe symptoms include >40 mmHg change in systolic blood pressure from baseline, increase in temperature with rigors, shortness of breath with wheezing, and / or stridor.

## Physician information

First Name: \_\_\_\_\_ Middle: \_\_\_\_\_ Last: \_\_\_\_\_ Practice: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ ZIP: \_\_\_\_\_

Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ NPI: \_\_\_\_\_ Contact: \_\_\_\_\_

By signing, I certify/recertify that the above therapy, products and services are medically necessary and that this patient is under my care. I have received authorization to release the above referenced information and medical and/or patient information relating to this therapy. Pharmacy has my permission to contact the insurance company on my behalf to obtain authorization for patient.

\_\_\_\_\_  
Substitution permissible signature

\_\_\_\_\_  
Dispense as written signature

\_\_\_\_\_  
Date

Please fax both pages of this completed form with a copy of any medical history and labs relevant to the prescribed therapy.

**This form is not a valid prescription in New York.**